WEST

annamanas (amamanamananamana)

Generate Collection

L4: Entry 4 of 16

File: DWPI

Oct 10, 1996

DERWENT-ACC-NO: 1996-464693

DERWENT-WEEK: 199646

COPYRIGHT 2001 DERWENT INFORMATION LTD

TITLE: Herbicidal compsn. contg. non-selective <u>herbicide</u> such as glyphosate - and adjuvant of organo-silicon, polyoxyethylene dodecyl ether, ethylene-oxide-propylene-oxide block copolymer, beef tallow amino-ether etc.

INVENTOR: HOSAKA, H; KOJIMA, S; YAMAGISHI, H

PRIORITY-DATA: 1995JP-0198189 (July 11, 1995), 1995JP-0106985 (April 6, 1995)

PATENT-FAMILY:

 PUB-NO
 PUB-DATE
 LANGUAGE
 PAGES
 MAIN-IPC

 WO 9631121 A1
 October 10, 1996
 J
 014
 A01N027/00

 JP 08333204 A
 December 17, 1996
 005
 A01N025/30

INT-CL (IPC): A01N 25/30; A01N 27/00; A01N 33/12; A01N 47/08; A01N 27/00; A01N 47/28

ABSTRACTED-PUB-NO: WO 9631121A BASIC-ABSTRACT:

Herbicidal compsn. comprises a nonselective <u>herbicide</u> and at least one adjuvant. The adjuvant is selected from organosilicon cpds., polyoxyethylene isotridecyl ether, polyethylene dodecyl ether, beef tallow amino ether, ethyleneoxy-propyleneoxy (EO/PO) block copolymers, EO/PO aliphatic alcohol ethers and polyoxyethylene nonylphenyl ethers.

Also claimed is a nonselective <u>herbicide</u>-contg. compsn. for use in an aq. spray that contains 0.01-10 wt.% the adjuvant.

USE - The compsn. may be used to treat <u>foliage</u> and soil, for killing of weeds in agriculture and horticulture.

ADVANTAGE - The compsn. provides rapid onset of herbicidal activity with improved herbicidal effect.

Generate Collection

L4: Entry 5 of 16

File: DWPI

Apr 7, 1993

DERWENT-ACC-NO: 1993-110918

DERWENT-WEEK: 199314

COPYRIGHT 2001 DERWENT INFORMATION LTD

TITLE: Herbicidal compsn. of glyphosate and a <u>silicone</u> adjuvant - has improved after-rain phytotoxicity, esp. in the control of velvet-leaf

INVENTOR: PETROFF, L J; SNOW, S A

PRIORITY-DATA: 1991US-0767960 (September 30, 1991)

PATENT-FAMILY:

PUB-NO		PUB-	PUB-DATE		GE PAGES	MAIN-IPC
	EP 535596	A1 Apri	1 7, 1993	E	011	A01N057/20
	AU 922539	9 A Apri	1 1, 1993		000	A01N025/24
	CA 207930	8 A Marc	h 31, 1993	3	000	A01N057/20
	JP 052389	10 A Sept	ember 17,	1993	008	A01N057/12

INT-CL (IPC): A01N 25/04; A01N 25/24; A01N 25/30; A01N 55/00; A01N 57/12; A01N 57/20; A01N 55/00; A01N 57/20

ABSTRACTED-PUB-NO: EP 535596A BASIC-ABSTRACT:

A compsn. comprises: foliar-applied N-phosphonomethylglycine (I) or its agriculturally acceptable salt; and 0.01 to 50 pts wt., per part of (I), of a silicone adjuvant (II) where x is 0 or 1; y is 1 or 2; and G is -R1N(+)(R2)3 X(-), -R1N(+)(R2)2(R3)-Y(-), -R1N(R2)2 or -R1y(-)M(+); in which R1 and R3 each represent a divalent 2-10C organic connecting group selected from alkylene, alkylene ether and alkylene thioether, each opt. OH-substd.; R2 is a 1-10C alkyl radical or a 1-5C hydroxyalkyl group; X(-) is Cl(-), Br(-), I(-), NO3(-), BF4(-), PF6(-), R4 is 1-5C alkyl; Y(-) is a terminal ionic group which is -COO(-) or -SO3(-); and M(+) is K(+) or Na(+).

USE/ADVANTAGE - The compsn. is useful as a herbicide, esp. against velvetleaf (Abutilon theophrasti), Application to the foliage is pref. by spraying at a rate which provides 0.5-10 pounds per acre of (I). The after-rain phytotoxicity of glyphosate herbicides is signficantly increased by the inclusion of silicone adjuvants (II), and this improvement in rain-fastness is obtd. without the need for an additional sticking agent. Smaller amts. of herbicide may be used to obtain the desired effect.

obtain the desired effect.

2 of 2



WEST

Generate Collection

L4: Entry 14 of 16

File: DWPI

Nov 2, 1984

DERWENT-ACC-NO: 1984-309680

DERWENT-WEEK: 198450

COPYRIGHT 2001 DERWENT INFORMATION LTD

TITLE: Selective <u>herbicide for foliage</u> treatment - contains 2-methyl delta'-isopr opyl valeryl anilide as active component

PRIORITY-DATA: 1983JP-0067773 (April 19, 1983)

PATENT-FAMILY:

PUB-NO PUB-DATE

LANGUAGE PAGES MAIN-IPC

JP 59193806 A November 2, 1984

004

INT-CL (IPC): A01N 37/22

ABSTRACTED-PUB-NO: JP59193806A

BASIC-ABSTRACT:

Herbicide contains 2-methyl-Delta' isopropylvaleryl anilide
(I).

- (I) can be pred. by reacting 4-isopropylaniline with 2-methyl-valeric acid chloride. The reaction is carried out in inert solvent such as benzene, toluene and xylene in presence of acid binder such as triethylamine, pyridine and potassium carbonate.
- (I) can be used alone or formulated as powder, granule, wettable powder, flowable, etc. Suitable solid carriers are clay, talc, calcium carbonate, diatomaceous earth, zeolite, silicon oxide, etc. Suitable liquid carriers are oils and organic solvents.

ADVANTAGE - (I) shows very broad herbicidal spectrum by <u>foliage</u> treatment, and no phytotoxicity to corn, wheat even at high concn. It can control almost all harmful weeds in upland.